

HOW PYROLYSIS

as Chemical Recycling Technology
Contributes to a Circular Carbon Economy
as Analysed by LCA

 **BASF**

We create chemistry

TOPIC

Significant amounts of mixed plastic waste (MPW) currently end up in incineration plants and landfills. It is important that such waste can be managed appropriately in line with circular economy principles. One emerging option for managing MPW is chemical recycling via pyrolysis. Pyrolysis processes aim at closing the loop by breaking down the plastics into their monomers or delivering gases and condensates which can be fed into the existing chemical processing streams. The research question is, how pyrolysis as chemical recycling technology contributes to a circular carbon economy.

METHOD

This study from BASF and Sphera uses Life Cycle Assessment (LCA) to compare the environmental impacts of pyrolysis of MPW with the established waste management alternatives in Germany. Three case studies are assessed to cover different perspectives for pyrolysis: to get rid of plastic waste and compare with other disposal methods like incineration or to produce chemical products and compare with their conventional virgin or mechanically recycled counterparts. For the latter perspective, the mass balance approach is applied. Pyrolysis oil is fed into the steam cracker at BASF where the primary output materials are ethylene and propylene. These are used to make numerous chemical products.

RESULT

With the pyrolysis technology, mixed plastic waste that is currently difficult to recycle can be utilized. Pyrolysis oil sourced from BASF partners has been used on pilot scale in a running BASF production network.

Primary data from a pyrolysis plant was collected and used for LCA modelling and result calculations. Overall, the results show – based on the made assumptions and the defined scope – that pyrolysis can serve as a high value waste management option regarding climate change and material efficiency. In two case studies, pyrolysis was shown to be the preferred option regarding climate change impact and resource use. Pyrolysis is preferred over incineration as an end-of-life treatment route of mixed plastic waste and is also preferred over fossil-based naphtha for production of virgin-grade plastic feedstock. The results show also that there is no significant difference in climate change results for recycled plastic products from pyrolysis and mechanical recycling of mixed plastic waste streams.

KEYWORDS

Circular carbon economy, mass balance approach, environmental evaluation by life cycle assessment (LCA), plastics waste, chemical recycling, pyrolysis



MORE INFORMATION AT
WWW.BASF.COM

**10TH INTERNATIONAL CONFERENCE
ON LIFE CYCLE MANAGEMENT**

September 5 – 8, 2021, Stuttgart/Germany

Authors:

CAROLIN DEREGOWSKI, BASF SE

✉ carolin.deregowski@basf.com

MAIKE HORLACHER, Sphera Solutions GmbH

✉ mhorlacher@sphera.com

CHRISTIAN KRUEGER, BASF SE

✉ christian.krueger@basf.com

PETER SALING, BASF SE

✉ peter.saling@basf.com